

Note to Readers: *EHP* strives to ensure that all journal content is accessible to all readers. However, some figures and Supplemental Material published in *EHP* articles may not conform to 508 standards due to the complexity of the information being presented. If you need assistance accessing journal content, please contact ehp508@niehs.nih.gov. Our staff will work with you to assess and meet your accessibility needs within 3 working days.

Supplemental Material

Effects of Laser Printer–Emitted Engineered Nanoparticles on Cytotoxicity, Chemokine Expression, Reactive Oxygen Species, DNA Methylation, and DNA Damage: A Comprehensive *in Vitro* Analysis in Human Small Airway Epithelial Cells, Macrophages, and Lymphoblasts

Sandra V. Pirela, Isabelle R. Miousse, Xiaoyan Lu, Vincent Castranova, Treye Thomas, Yong Qian, Dhimiter Bello, Lester Kobzik, Igor Koturbash, and Philip Demokritou

Table of Contents

Supplemental Material, Part A

Dosimetric considerations for *in vitro* testing – Example of calculations

Table S1. Summary of parameters used in the *in vivo* lung Multiple Path Particle Deposition model (MPPD2).

Table S2. Assays for determination of LINE-1 and *Alu* methylation.

Table S3. Assays for determination of gene expression.

Table S4. *In vitro* administered and delivered doses of SiO₂ and MS-WF.

Figure S1. Hydrodynamic diameter as a function of DSE for PEPs (PM_{0.1}) and MS-WF. DSE_{cr}: critical delivered sonication energy, energy required for minimal agglomeration.

Figure S2. Fraction of administered dose deposited, f_D , as a function of *in vitro* exposure time for PEPs (PM_{0.1}), SiO₂ and MS-WF calculated using the agglomeration diameter and estimated effective density. Plots are presented for the tested materials in the two media formulations (RPMI/10% FBS and SAGM).

Figure S3. Deposition mass flux (left axis) and deposition fraction (right axis) as a function of airway generation number.

Figure S4. Quantitative DNA damage assessment (Comet assay) of human lymphoblasts (TK6 cells) exposed for 4 hours to PEPs at various doses. All values are represented as mean \pm SE.

References for Supplemental Material, Part A.